

Quantum Knizhnik-Zamolodchikov equations and holomorphic vector bundles¹

Pavel I. Etingof

Yale University
Department of Mathematics
2155 Yale Station
New Haven, CT 06520 USA

e-mail etingof@ pascal.math.yale.edu

July 1992

Introduction

In 1984 Knizhnik and Zamolodchikov [KZ] studied the matrix elements of intertwining operators between certain representations of affine Lie algebras and found that they satisfy a holonomic system of differential equations which are now called the Knizhnik-Zamolodchikov (KZ) equations. It turned out that the KZ equations (and hence, representation theory of affine Lie algebras) are a source of a rich variety of special functions. The Gauss hypergeometric function and its various generalizations were obtained as solutions of the KZ system.

In the recent paper [FR], Frenkel and Reshetikhin considered intertwining operators between representations of quantum affine algebras. It was shown that their matrix elements satisfy a system of holonomic difference equations – a natural quantum analogue of the KZ system. It was also shown that the solutions of this difference system are very nontrivial special functions that generalize basic hypergeometric series. In particular, one of the simplest solutions is expressed in terms of the q -hypergeometric function which was introduced by Heine in the 19th century. This is consistent with the general idea that all reasonable special functions should come from representation theory, as matrix elements of certain representations.

The classical KZ system can be regarded as a local system – a flat structure in the trivial vector bundle on the space of sets of N distinct points on the complex plane. This fact enabled Schechtman and Varchenko to apply geometric methods to the study of the KZ equations and obtain a complete solution for a general simple Lie algebra [SV]. This interpretation can be extended to the quantum case [M;R] with the help of a specially designed discrete analogue of the notion of a local system. Other than that, the geometric meaning of the quantum KZ equations has so far remained unclear.

The goal of this paper is to introduce a new geometric interpretation of the quantum KZ equations. In Chapters 1 and 2, they are linked to certain holomorphic vector bundles on a product of N equivalent elliptic curves, naturally constructed by a gluing procedure from a system of trigonometric quantum R -matrices. Meromorphic solutions of the quantum KZ equations are interpreted as sections of such a bundle. This interpretation is an analogue of the interpretation of solutions of the classical KZ equations as sections of a flat vector bundle. It yields a non-technical proof of consistency of the quantum KZ system.

In Chapter 3 it is shown that the matrix elements of intertwiners between representations of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ correspond to regular (holomorphic) sections.

In Chapter 4, Birkhoff's connection matrices for the quantum KZ equations are interpreted as transition matrices from one fundamental system of holomorphic sections to another. They can be used to give an alternative construction of the holomorphic vector bundle corresponding to the quantum KZ system.

In Chapter 5, the topological structure of the vector bundle associated to the

quantum KZ equations is studied in the special case of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}_2})$. The Chern classes of this bundle are evaluated, and it is shown that they uniquely determine its topology. The main tool of this evaluation is the crystal limit $q \rightarrow 0$, and its result shows that the topology of the quantum KZ equations encodes the structure of the crystal bases in representations of the quantum group $U_q(\mathfrak{sl}_2)$.

Chapter 6 treats the special case $N = 2$. In this case, one essentially deals with a vector bundle over an elliptic curve. This bundle is shown to be semistable (for the case of $U_q(\widehat{\mathfrak{sl}_2})$) if the parameters take generic values. The proof makes use of the crystal limit $q \rightarrow 0$.

In Chapter 7, we give a vector bundle interpretation of the generalized quantum KZ equations for arbitrary affine root systems defined recently by Cherednik [Ch].

For the sake of brevity, the results regarding quantum groups are formulated and proved for $U_q(\mathfrak{sl}_2)$. Most of them can be suitably extended to $U_q(\mathfrak{g})$ where $\mathfrak{g}$ is an arbitrary simple Lie algebra.

Acknowledgements

This paper was inspired by the course on the classical and quantum Knizhnik-Zamolodchikov equations given by my adviser Igor Frenkel at Yale in the spring of 1992. It is a pleasure for me to thank Professor Frenkel for helping me build up a necessary background and guiding my work on this paper.

I would also like to thank Professors David Kazhdan, Nikolai Reshetikhin, and Alexandre Varchenko for useful discussions.

1. Trigonometric R -matrices and holomorphic vector bundles.

Let $V_1, \dots, V_N$ be a collection of finite-dimensional vector spaces. Let $W = V_1 \otimes V_2 \otimes \dots \otimes V_N$.

Let $R \in \text{End}(V_i \otimes V_j)$, $R = \sum_n a_n \otimes b_n$, $a_n \in \text{End}(V_i)$, $b_n \in \text{End}(V_j)$. Throughout the paper we will use the same notation R for the operator $\sum_n 1_{U_1} \otimes a_n \otimes 1_{U_2} \otimes b_n \otimes 1_{U_3}$ in $\text{End}(U_1 \otimes V_i \otimes U_2 \otimes V_j \otimes U_3)$ and the operator $\sum_n 1_{U_1} \otimes b_n \otimes 1_{U_2} \otimes a_n \otimes 1_{U_3}$ in $\text{End}(U_1 \otimes V_j \otimes U_2 \otimes V_i \otimes U_3)$, where U_1, U_2, U_3 are arbitrary vector spaces and $1_U \in \text{End}(U)$ is the identity map.

Definition 1.1. *A system of trigonometric quantum R -matrices on $V_1, \dots, V_N$ is a collection of rational functions $\{R_{ij}(z)\}$ with values in $\text{End}(V_i \otimes V_j)$ for $1 \leq i, j \leq N$, $i \neq j$ such that*

(i) $R_{ij}(z)$ satisfy the quantum Yang-Baxter equation:

$$(1.1) \quad R_{ij} \left(\frac{z_1}{z_2} \right) R_{ik} \left(\frac{z_1}{z_3} \right) R_{jk} \left(\frac{z_2}{z_3} \right) = R_{jk} \left(\frac{z_2}{z_3} \right) R_{ik} \left(\frac{z_1}{z_3} \right) R_{ij} \left(\frac{z_1}{z_2} \right)$$

in $\text{End}(W)$, $1 \leq i, j, k \leq N$;

(ii) $R_{ij}(z)$ satisfy the unitarity condition

$$(1.2) \quad R_{ij}(z) = R_{ji}(z^{-1})^{-1}$$

in $\text{End}(W)$, $1 \leq i, j \leq N$;

(iii) $R_{ij}(z)$ have no poles on the unit circle.

Definition 1.2. *A collection of linear transformations $A_i \in GL(W)$, $1 \leq i \leq N$, is compatible to the system of matrices $\{R_{ij}(z)\}$ if*

$$(1.3) \quad \begin{aligned} A_i R_{jk}(z) &= R_{jk}(z) A_i, \quad i \neq j, k, \\ A_i A_j R_{ij}(z) &= R_{ij}(z) A_j A_i, \quad 1 \leq i, j \leq N \end{aligned}$$

in $\text{End}(W)$.

Let $p \in \mathbb{C}^*$, $|p| < 1$, and let Π be the subgroup of $\mathbb{C}^*$ generated by p . Consider the complex torus $T = \mathbb{C}^* / \Pi$. We will realize T as an annulus $\{z \in \mathbb{C}^* \mid |p| \leq |z| < 1\}$ with identified boundaries. Denote by T^N the direct product of N copies of T .

It turns out that to any system of trigonometric quantum R -matrices on $V_1, \dots, V_N$ and any collection of linear transformations compatible to this system one can naturally associate a holomorphic vector bundle on T^N with fiber W .

Partition T^N into $N!$ chambers $D_s = \{(z_1, \dots, z_N) \mid |p| < |z_{s(N)}| < \dots < |z_{s(1)}| < 1\}$, where s is a permutation from the symmetric group S_N . These sets are open and disjoint, and their closures cover T^N . Also, they are permuted by the natural action of S_N on T^N : $s_1(D_{s_2}) = D_{s_2 s_1}$, where $s(z_1, \dots, z_N) = (z_{s(1)}, \dots, z_{s(N)})$.

We say that two chambers D_{s_1} and D_{s_2} are adjacent if $\partial D_{s_1} \cap \partial D_{s_2}$ is a hypersurface of real codimension 1 in T^N . The following trivial lemma classifies pairs of adjacent chambers.

Lemma 1.1. *Chambers D_{s_1} and D_{s_2} are adjacent if and only if $s_1^{-1} s_2$ is either a transposition of adjacent elements $t_j = (j, j+1)$, $1 \leq j \leq N-1$, or a cyclic permutation $c^{\pm 1}$, $c = (N, N-1, \dots, 2, 1)$.*

The entire torus can be obtained by gluing the chambers together along the boundaries between them. Therefore, in order to define a vector bundle on T^N

it suffices to proclaim it trivial over each of the chambers and prescribe transition matrices on the boundaries between adjacent chambers. These boundaries are

$$\Gamma_{s,j} = \{(z_1, \dots, z_N) \mid |p| < |z_{s(N)}| < \dots < |z_{s(j+1)}| = |z_{s(j)}| < \dots < |z_{s(1)}| < 1\}$$

and

$$\Sigma_s = \{(z_1, \dots, z_N) \mid |p| = |z_{s(N)}| < \dots < |z_{s(1)}| < 1\}.$$

Note that $\Gamma_{s,j} = \Gamma_{st_j,j}$.

Now let $\{R_{ij}(z)\}$ be any system of rational functions with values in $\text{End}(V_i \otimes V_j)$ satisfying condition (iii), and let $\{A_i\}$ be a system of linear transformations compatible to $\{R_{ij}(z)\}$.

On $\Gamma_{s,j}$, define the transition matrices from D_s to D_{st_j} to be

$$(1.4) \quad T_{s,j}(z_1, \dots, z_N) = R_{s(j+1)s(j)} \left(\frac{z_{s(j+1)}}{z_{s(j)}} \right)$$

On Σ_s , set the transition matrices from D_s to D_{sc} to be

$$(1.5) \quad Q_s = A_{s(N)}$$

These transition matrices take values in $\text{End}(W)$.

Theorem 1.2. *Transition matrices (4) and (5) define a holomorphic vector bundle on T^N with fiber W if and only if $\{R_{ij}(z)\}$ is a system of trigonometric R -matrices and $\{A_i\}$ are compatible to $\{R_{ij}(z)\}$.*

Remark 1.1. This fact can be considered as a geometric interpretation of trigonometric solutions to the quantum Yang-Baxter equation with a complex parameter.

Proof. First of all, the regularity of $\{R_{ij}(z)\}$ on the unit circle is equivalent to the smoothness of the transition functions on the boundaries between chambers. Thus, it remains to show that the consistency conditions on transition matrices reduce to relations (1.1), (1.2) and (1.3).

The consistency conditions apply at every point $P \in T^N$ whose arbitrarily small neighborhood intersects more than two chambers. Let U be a small enough neighborhood of P . Consider the graph $G(P)$ whose vertices are connected components of intersections of chambers with U .² Two vertices are connected with an edge if the corresponding two components are adjacent inside U . Thus every edge is labeled with a transition matrix – a holomorphic $\text{End}(W)$ -valued function in U . Hence, a holomorphic in U transition matrix can be associated to every oriented path in the graph $G(P)$. This matrix is defined to be the product of the transition matrices labeling the edges of this path in the opposite order to that prescribed by the orientation. The consistency conditions demand that any two paths with the same beginning and end be labeled with the same transition matrix (this is equivalent to the uniqueness of analytic continuation of sections of the bundle in U).

Let $P = (z_1, \dots, z_N)$. First assume that $|z_j| \neq |p|$ for all j . Let s be a permutation labeling one of the chambers intersecting with U . Then there exists an increasing sequence $1 \leq j_1 < j_2 < \dots < j_r = N$ such that

$$(1.6) \quad |z_{s(1)}| = \dots = |z_{s(j_1)}| > |z_{s(j_1+1)}| = \dots = |z_{s(j_2)}| > \dots > |z_{s(j_{r-1}+1)}| = \dots = |z_{s(j_r)}|.$$

²Note that although the chambers are connected, their intersections with U may consist of

Obviously, vertices of $G(P)$ correspond to permutations of the form $s\sigma$, $\sigma \in S(P)$, where $S(P) = S_{j_1} \times S_{j_2-j_1} \times \cdots \times S_{j_r-j_{r-1}} \subset S_N$. Denote such a vertex by $v(\sigma)$. It is clear that $v(\sigma_1)$ and $v(\sigma_2)$ are connected with an edge if and only if $\sigma_1 = \sigma_2 t_j$, $j \neq j_1, \dots, j_{r-1}$. It follows that a path in $G(P)$ is a representation of an element $\sigma \in S(P)$ as a product of the generators t_j . It is known from the theory of the symmetric group that any two such representations can be identified with the help of relations

$$(1.7) \quad t_j t_{j+1} t_j = t_{j+1} t_j t_{j+1},$$

$$(1.8) \quad t_j^2 = 1,$$

$$(1.9) \quad t_i t_j = t_j t_i, \quad |i - j| > 1.$$

Applying this result to the transition matrices, we find that the consistency conditions at P are equivalent to identities (1.1) and (1.2) for the matrices $\{R_{ij}(z)\}$. Indeed, it is easy to see that relation (1.7) reduces to the quantum Yang-Baxter equation, relation (1.8) yields the unitarity, and relation (1.9) trivially follows from the definition of transition matrices.

Now assume that (1.6) holds and $|z_{s(j)}| = |p|$ for $j_{r-1} < j \leq j_r$. Set $k = j_r - j_{r-1}$. The graph $G(P)$ can then be represented as a union $G(P) = G_0(P) \cup G_1(P) \cup \cdots \cup G_k(P)$, where $G_l(P)$ corresponds to the points in the vicinity of U with $|z_{s(j)}|$ close to $|p|$ for $j_{r-1} < j \leq j_r - l$ and close to 1 for $j_r - l < j \leq j_r$. Vertices of $G_l(P)$ are labeled by permutations from $S(P)$ as follows: $v_l(\sigma) \in G_l(P)$ corresponds to the connected component of $D_{s\sigma c^l}$. Two vertices $v_l(\sigma_1)$ and $v_l(\sigma_2)$ in $G_l(P)$ are connected with an edge if and only if $\sigma_1 c^l = \sigma_2 c^l t_j$, $j \neq l, j_1 + l, \dots, j_{r-1} + l$. Also, two vertices $v_l(\sigma_1) \in G_l(P)$ and $v_m(\sigma_2) \in G_m(P)$, $l \neq m$, are connected if and only if $|l - m| = 1$ and $\sigma_1 = \sigma_2$. This completely describes the structure of the graph $G(P)$.

Let $k \geq 1$. For $l \leq k - 1$ consider the paths $\pi_1(\sigma, l) : v_l(\sigma) \rightarrow v_l(\sigma t_j) \rightarrow v_{l+1}(\sigma t_j)$ and $\pi_2(\sigma, l) : v_l(\sigma) \rightarrow v_{l+1}(\sigma) \rightarrow v_{l+1}(\sigma t_j)$, $j \neq j_1, \dots, j_{r-1}, j_r - l, j_r - l - 1$. These paths have a common beginning and ending, and the fact that they give the same transition matrix is expressed by the relation $R_{\sigma(j+1)\sigma(j)} A_{\sigma(N-l)} = A_{\sigma(N-l)} R_{\sigma(j+1)\sigma(j)}$ which is the first part of (1.3).

Now let $k \geq 2$, and for $l \leq k - 2$ consider the paths

$$\pi_1^1(\sigma, l) : v_l(\sigma) \rightarrow v_l(\sigma t_{N-l-1}) \rightarrow v_{l+1}(\sigma t_{N-l-1}) \rightarrow v_{l+2}(\sigma t_{N-l-1}) \text{ and}$$

$$\pi_2^1(\sigma, l) : v_l(\sigma) \rightarrow v_{l+1}(\sigma) \rightarrow v_{l+2}(\sigma) \rightarrow v_{l+2}(\sigma t_{N-l-1}),$$

with the same beginning and ending. The fact that they should give the same transition matrix is expressed by the relation

$$R_{\sigma(N-l)\sigma(N-l-1)} A_{\sigma(N-l-1)} A_{\sigma(N-l)} = A_{\sigma(N-l)} A_{\sigma(N-l-1)} R_{\sigma(N-l)\sigma(N-l-1)},$$

which is the second part of (1.3).

An elementary combinatorial argument shows that any two paths in the graph $G(P)$ with the same beginning and ending can be identified with each other by replacing subpaths of the form π_1 by π_2 , π_1^1 by π_2^1 , and vice versa, and using relations (1.7)–(1.9) when necessary. This means that once (1.1)–(1.3) are satisfied, any two such paths are forced to have the same transition matrix. Thus, we have shown that the consistency conditions on transition matrices are equivalent to relations (1.1), (1.2), and (1.3). ■

From now on we assume that the matrices $\{R_{ij}(z)\}$ and $\{A_i\}$ are fixed and denote the vector bundle constructed above by E_N .

Let $\Delta = (\Delta_1, \dots, \Delta_N)$ be a set of complex numbers. Introduce a new holomorphic vector bundle $E_\Delta(\Delta)$ obtained from E_N by twisting. For $\delta \in \mathbb{C}$ define a line

bundle L_δ on T whose meromorphic sections are functions $\psi(z)$ meromorphic in $\mathbb{C}^*$ such that $\psi(pz) = p^\delta \psi(z)$. Clearly, the isomorphism class of L_δ is determined by $\delta \bmod(1, 2\pi\sqrt{-1}/\log p)$, or, equivalently, by the image of $p^\delta \in \mathbb{C}^*$ in T . Let $\theta_j : T^N \rightarrow T$, $1 \leq j \leq N$ be the projection: $\theta_j(z_1, \dots, z_N) = z_j$, and let $\mathcal{B}_j(\Delta_j) = \theta_j^*(L_{\Delta_j})$. $\mathcal{B}_j(\Delta_j)$ are topologically trivial holomorphic line bundles on T^N . Let $\mathcal{B}(\Delta) = \mathcal{B}_1(\Delta_1) \otimes \mathcal{B}_2(\Delta_2) \otimes \dots \otimes \mathcal{B}_N(\Delta_N)$. Now for any bundle $\mathcal{E}$ over T^N define a new bundle $\mathcal{E}(\Delta)$ by

$$(1.10) \quad \mathcal{E}(\Delta) = \mathcal{E} \otimes \mathcal{B}(\Delta).$$

In the next section we will identify solutions of the quantum KZ equations with sections of $E_N(\Delta)$.

2. Quantum Knizhnik-Zamolodchikov equations and meromorphic sections.

Let $C = \{(z_1, \dots, z_N) \in T^N : 1 > |z_j| > |p|, 1 \leq j \leq N\}$. Let $\phi(z_1, \dots, z_N)$ be a meromorphic function in $\mathbb{C}^{*N}$ with values in W . Set $\psi(z_1, \dots, z_N) = z_1^{\Delta_1} \dots z_N^{\Delta_N} \phi(z_1, \dots, z_N)$.

To the function ψ we can associate a meromorphic section of $E_N(\Delta)$ over the open set C as follows.

A meromorphic section of $E_N(\Delta)$ over C is a collection of functions $\psi_s(z_1, \dots, z_N) = z_1^{\Delta_1} \dots z_N^{\Delta_N} \phi_s(z_1, \dots, z_N)$, $s \in S_N$, where ϕ_s is a W -valued meromorphic function on $D_s^c \cap C$ (D_s^c denotes the closure of the chamber D_s), satisfying the consistency conditions

$$(2.1) \quad \phi_{st_j} = T_{s,j} \phi_s \text{ on } \Gamma_{s,j}.$$

Let w be the element of maximal length in S_N ($w(i) = N - i + 1$). To construct a section of $E_N(\Delta)$ from ψ , set $\phi_w = \phi$, and then extend it to the whole set C by applying rule (2.1). That means, if $s = t_{j_1} \dots t_{j_n}$ and $s_l = t_{j_1} \dots t_{j_l}$, $1 \leq l \leq n$, then set

$$(2.2) \quad \phi_{ws} = T_{ws_{n-1}, j_n} T_{ws_{n-2}, j_{n-1}} \dots T_{ws_1, j_2} T_{w, j_1} \phi$$

in D_{ws} . It follows from relations (1.1) and (1.2) that the result of this extension does not depend on the decomposition of s in the product of t_j . Clearly, functions (2.2) satisfy conditions (2.1). Thus, we have constructed a section of the bundle $E_N(\Delta)$ over C . Denote this section by $\tilde{\psi}$.

Now let us define the quantum KZ equations – the main subject of this paper.

Definition 2.1. *The difference equations on a W -valued function ψ*

$$(2.3) \quad \begin{aligned} \psi(z_1, \dots, pz_j, \dots, z_N) &= R_{j,j-1} \left(\frac{z_j}{z_{j-1}} p \right) R_{j,j-2} \left(\frac{z_j}{z_{j-2}} p \right) \dots R_{j,1} \left(\frac{z_j}{z_1} p \right) \times \\ &A_j R_{j,N} \left(\frac{z_j}{z_N} \right) R_{j,N-1} \left(\frac{z_j}{z_{N-1}} \right) \dots R_{j,j+1} \left(\frac{z_j}{z_{j+1}} \right) \psi(z_1, \dots, z_j, \dots, z_N), \quad 1 \leq j \leq N, \end{aligned}$$

are called the quantum Knizhnik-Zamolodchikov equations.

The following theorem gives a new geometric interpretation of the quantum KZ equations

Theorem 2.1. *The section $\tilde{\psi}$ of the bundle $E_N(\Delta)$ over C extends to a global meromorphic section of this bundle if and only if the function $\psi(z_1, \dots, z_N)$ satisfies the quantum KZ equations.*

Proof. In order for $\tilde{\psi}$ to extend, it is necessary and sufficient that ψ_s defined above satisfy the additional consistency conditions on the surfaces Σ_s . Obviously, it is enough to require consistency only for $s = wt_j t_{j+1} \dots t_{N-1}$, $1 \leq j \leq N$. In this case, the conditions are

$$(2.4) \quad \phi_{sc}(z_1, \dots, pz_j, \dots, z_N) = A_j \phi_s(z_1, \dots, z_j, \dots, z_N).$$

Using the decomposition $c = t_{N-1} t_{N-2} \dots t_1$ and equations (2.2) and (1.4), we obtain from (2.4)

$$(2.5) \quad R_{1,j} \left(\frac{z_1}{z_j p} \right) \dots R_{j-2,j} \left(\frac{z_{j-2}}{z_j p} \right) R_{j-1,j} \left(\frac{z_{j-1}}{z_j p} \right) \psi(z_1, \dots, pz_j, \dots, z_N) = \\ A_j R_{j,N} \left(\frac{z_j}{z_N} \right) R_{j,N-1} \left(\frac{z_j}{z_{N-1}} \right) \dots R_{j,j+1} \left(\frac{z_j}{z_{j+1}} \right) \psi(z_1, \dots, z_j, \dots, z_N),$$

which is equivalent to (2.3). ■

Corollary 2.2. *The quantum KZ equations are consistent.*

Proof. Any holomorphic vector bundle on a compact complex manifold has nonzero meromorphic sections. Therefore, equations (2.3) have nonzero solutions, which implies that they are consistent. ■

Definition 2.2. *Let us say that a system of trigonometric R -matrices $\{R_{ij}(z)\}$ is regular if for every pair i, j such that $i \neq j$*

- (i) *$R_{ij}(z)$ is regular and nondegenerate at the origin and infinity;*
- (ii) *$R_{ij}(z)$ is regular outside the unit circle and $R_{ij}^{-1}(z)$ is regular inside the unit circle.*

From now on we assume that $\{R_{ij}(z)\}$ is regular and use the notation

$$(2.6) \quad M_j(z_1, \dots, z_N) = \\ R_{j,j-1} \left(\frac{z_j}{z_{j-1} p} \right) \dots R_{j,1} \left(\frac{z_j}{z_1 p} \right) A_j R_{j,N} \left(\frac{z_j}{z_N} \right) \dots R_{j,j+1} \left(\frac{z_j}{z_{j+1}} \right)$$

Fix $s \in S_N$. Let

$$(2.7) \quad M_j^s = \lim_{z_{s(i)}/z_{s(i+1)} \rightarrow \infty, 1 \leq i \leq N-1} M_j(z_1, \dots, z_N).$$

It follows from the consistency of the quantum KZ equations that

$$(2.8) \quad [M_i^s, M_j^s] = 0$$

for any pair i, j .

Theorem 2.3. [FR]

(i) There exists a matrix solution of the quantum KZ equations of the form

$$(2.9) \quad L_s(z_1, \dots, z_N) = z_1^{\frac{\log M_1^s}{\log p}} \dots z_N^{\frac{\log M_N^s}{\log p}} F_s(z_1, \dots, z_N),$$

such that F_s is an $\text{End}(W)$ -valued meromorphic function in $\mathbb{C}^{*N}$ regular in the region $|z_{s(1)}| > |z_{s(2)}| > \dots > |z_{s(N)}|$ with $\lim_{z_{s(i)}/z_{s(i+1)} \rightarrow \infty, 1 \leq i \leq N-1} F_s = 1_W$ (this function will be homogeneous of degree 0).

(ii) Any vector solution ψ of the quantum KZ equations regular in the region $|z_{s(1)}| > |z_{s(2)}| > \dots > |z_{s(N)}|$ has the form $L_s u$ where $u \in W$.

Idea of proof. The solution L_s is given by the following limit:

$$(2.10) \quad \lim_{k_{s(j_1)} - k_{s(j_2)} \rightarrow \infty, j_1 > j_2} \prod_{j=1}^N \prod_{i=0}^{k_j-1} M_j^{-1}(z_1 p^{k_1}, \dots, z_{j-1} p^{k_{j-1}}, z_j p^i, z_{j+1}, \dots, z_N) \prod_{j=1}^N (M_j^s)^{k_j}.$$

The existence of this limit follows from the results of [Ao].

Theorem 2.4. Let ϕ and ψ be as above. If $\{R_{ij}(z)\}$ is regular and $\phi(z_1, \dots, z_N)$ is holomorphic in the region $1 \geq |z_{s(1)}| \geq |z_{s(2)}| \geq \dots \geq |z_{s(N)}| \geq |p|$, $s \in S_N$, then $\tilde{\psi}$ is a holomorphic section of the bundle $E_N(\Delta)$.

Proof. We will assume that $s = id$. For an arbitrary permutation s , the proof is similar.

First of all, let us show that $\phi(z_1, \dots, z_N)$ is holomorphic in the annulus $\mathcal{A} = \{(z_1, \dots, z_N) \in \mathbb{C}^N : |p| \leq |z_1|, \dots, |z_N| \leq 1\}$. Let J be an integer such that all $R_{ij}(z)$ are holomorphic whenever $|z| \leq |p|^J$. We will use the notation $\hat{z}_j = p^{J(j-1)} z_j$. Since ψ satisfies the quantum KZ equations, we have

$$(2.11) \quad \phi(z_1, \dots, z_N) = \text{const} \prod_{j=2}^N \prod_{i=0}^{(j-1)J-1} M_j^{-1}(z_1, \dots, z_{j-1}, p^i z_j, \hat{z}_{j+1}, \dots, \hat{z}_N) \phi(\hat{z}_1, \dots, \hat{z}_N)$$

Since $\{R_{ij}(z)\}$ is regular, $M_j^{-1}(z_1, \dots, z_N)$ is holomorphic if $z_j p \leq z_i$ for $i < j$ and $z_j \leq z_i$ for $i > j$. This implies that all the factors M_j^{-1} in (2.11) are holomorphic in $z_1, \dots, z_N$ in $\mathcal{A}$. The function $\phi(\hat{z}_1, \dots, \hat{z}_N)$ is also holomorphic in the annulus, since $|\hat{z}_1| < \dots < |\hat{z}_N|$ whenever $(z_1, \dots, z_N) \in \mathcal{A}$. This shows that $\phi(z_1, \dots, z_N)$ is holomorphic in the annulus.

Now we are in a position to prove the holomorphicity of the section $\tilde{\psi}$. Since ϕ is holomorphic in $\mathcal{A}$, clearly $\tilde{\psi}$ is holomorphic in D_w . Let us analytically continue $\tilde{\psi}$ into the region D_{ws} , $s \in S_N$, along some path. Let $s = t_{m_1} \dots t_{m_l}$ be a minimal length decomposition of s . Then over D_{ws} $\tilde{\psi}$ is represented by a function of the form

$$(2.12) \quad \psi_{ws}(z_1, \dots, z_N) = R_{i_l j_l} \left(\frac{z_{i_l}}{z_{j_l}} \right) \dots R_{i_1 j_1} \left(\frac{z_{i_1}}{z_{j_1}} \right) \psi(z_1, \dots, z_N).$$

The key property of this decomposition is that $|z_{i_m}|$ is always greater than $|z_{j_m}|$ in D_w . It follows from the fact that the decomposition of s we used had minimal

length. This property and the regularity of the R -matrices imply that ψ_{ws} is a holomorphic (multivalued) function in D_{ws} which proves that $\tilde{\psi}$ is a holomorphic section of $E_N(\Delta)$. ■

Let $s \in S_N$ be a permutation. Let $d = \dim W$, and let $u_1^s, \dots, u_d^s$ be the basis of W such that $M_j^s u_i = \Delta_j^{(s,i)} u_i$, $1 \leq j \leq N$, $1 \leq i \leq d$ (we assume the generic situation when such a basis exists). Let

$$(2.13) \quad \psi^{(s,i)} = L_s u_i$$

, and let $\tilde{\psi}^{(s,i)}$ be the corresponding holomorphic sections of $E_N(\Delta^{(s,i)})$.

The following proposition is a corollary of formula (2.10).

Proposition 2.5. *The sections $\tilde{\psi}^{(s,i)}(z_1, \dots, z_N)$ form a basis of the fiber W everywhere except points where $\frac{z_{s(j_1)}}{z_{s(j_2)}} p^n$ is a pole of $R_{s(j_1)s(j_2)}(z)$ for a suitable $n \in \mathbb{Z}$ and pair of indices j_1, j_2 such that $j_1 > j_2$. At such points, $\tilde{\psi}^{(s,i)}(z_1, \dots, z_N)$ are linearly dependent.*

Remark 2.1. We can legitimately talk about linear dependence or independence of $\tilde{\psi}^{(s,i)}$ despite they are sections of different bundles, because the projectivizations of all these bundles are isomorphic to each other.

Remark 2.2. Proposition 2.5 shows that $\tilde{\psi}^{(s,i)}$ play the role of a fundamental system of sections: they give a coordinate frame of the fiber at all points of the base except those lying on a finite set of hypersurfaces in T^N of complex codimension 1.

3. Matrix elements of intertwining operators for $U_q(\widehat{\mathfrak{sl}_2})$ and holomorphic sections.

The quantum affine algebra $U_q(\widehat{\mathfrak{sl}_2})$ is a Hopf algebra obtained by a standard q -deformation of the universal enveloping algebra of the Kac-Moody algebra $\widehat{\mathfrak{sl}_2}$ [D;J1]. As an associative algebra, it is generated by elements $e_i, f_i, K_i^{\pm 1}, i = 0, 1$ which satisfy the following relations:

$$(3.1) \quad \begin{aligned} e_i f_i - f_i e_i &= \frac{K_i - K_i^{-1}}{q - q^{-1}}, \\ e_i f_j - f_j e_i &= 0, i \neq j, \\ K_i e_i K_i^{-1} &= q^2 e_i, \quad K_i e_j K_i^{-1} = q^{-2} e_j, \quad i \neq j \\ K_i f_i K_i^{-1} &= q^{-2} f_i, \quad K_i f_j K_i^{-1} = q^2 f_j, \quad i \neq j \\ K_i K_j - K_j K_i &= 0, \\ e_i^3 e_j - \frac{q^3 - q^{-3}}{q - q^{-1}} e_i^2 e_j e_i + \frac{q^3 - q^{-3}}{q - q^{-1}} e_i e_j e_i^2 - e_j e_i^3 &= 0, i \neq j, \\ f_i^3 f_j - \frac{q^3 - q^{-3}}{q - q^{-1}} f_i^2 f_j f_i + \frac{q^3 - q^{-3}}{q - q^{-1}} f_i f_j f_i^2 - f_j f_i^3 &= 0, i \neq j. \end{aligned}$$

The comultiplication in $U_q(\widehat{\mathfrak{sl}_2})$ is defined by

$$(3.2) \quad \begin{aligned} \Delta(K_i) &= K_i \otimes K_i, \\ \Delta(e_i) &= e_i \otimes K_i + 1 \otimes e_i, \\ \Delta(f_i) &= f_i \otimes 1 + K_i^{-1} \otimes f_i \end{aligned}$$

and the antipode acts according to

$$(3.3) \quad S(e_i) = -e_i K_i^{-1}, \quad S(f_i) = -K_i f_i, \quad S(K_i) = K_i^{-1}.$$

Here q is a complex number. We will assume that q is not 0 and not a root of unity.

The algebra $U_q(\widehat{\mathfrak{sl}_2})$ can be extended by adding elements $D^{\pm 1}$ which satisfy the relations

$$(3.4) \quad DK_i = K_i D, \quad De_0 = qe_0 D, \quad Df_0 = q^{-1}f_0 D, \quad De_1 = e_1 D, \quad Df_1 = f_1 D$$

The algebra obtained by this extension is denoted by $U_q(\widetilde{\mathfrak{sl}_2})$.

Two kinds of representations are defined for $U_q(\widetilde{\mathfrak{sl}_2})$: Verma modules $V_{\lambda, k}$ and finite dimensional evaluation representations $V_\mu(z)$.

The Verma module $V_{\lambda, k}$ is generated by a highest weight vector v satisfying the relations

$$(3.5) \quad e_i v = 0, \quad K_1 v = q^\lambda v, \quad K_0 v = q^{k-\lambda} v, \quad k, \lambda \in \mathbb{C}.$$

This module is free over the subalgebra generated by $\widehat{f_i}$ and should be regarded as a quantum deformation of the Verma module over $\widehat{\mathfrak{sl}_2}$.

The evaluation representation $V_\mu(z)$ is defined with the help of the quantum group $U_q(\mathfrak{sl}_2)$ (see e.g. [CP]). $U_q(\mathfrak{sl}_2)$ is a Hopf algebra generated by elements $e, f, K^{\pm 1}$ satisfying the relations

$$(3.6) \quad \begin{aligned} ef - fe &= \frac{K - K^{-1}}{q - q^{-1}}, \\ KeK^{-1} &= q^2 e, \\ KfK^{-1} &= q^{-2} f, \end{aligned}$$

in which the comultiplication and the antipode are given by

$$(3.7) \quad \begin{aligned} \Delta(K) &= K \otimes K, \\ \Delta(e) &= e \otimes K + 1 \otimes e, \\ \Delta(f) &= f \otimes 1 + K^{-1} \otimes f, \end{aligned}$$

$$(3.8) \quad \begin{aligned} S(e) &= -eK^{-1}, \\ S(f) &= -Kf, \\ S(K) &= K^{-1}. \end{aligned}$$

For $z \in \mathbb{C}^*$ Jimbo J2 defined the canonical algebra homomorphisms

$$p_z : U_q(\widehat{\mathfrak{sl}_2}) \rightarrow U_q(\mathfrak{sl}_2)$$

as follows:

$$(3.9) \quad \begin{aligned} p_z(e_1) &= e, \quad p_z(f_1) = f, \quad p_z(K_1^{\pm 1}) = K^{\pm 1}, \\ p_z(e_0) &= qf, \quad p_z(f_0) = q^{-1}e, \quad p_z(K^{\pm 1}) = K^{\mp 1} \end{aligned}$$

Let V_μ be the finite dimensional irreducible representation of $U_q(\mathfrak{sl}_2)$ with the highest weight μ – a nonnegative integer. Such a representation is unique for any μ . The homomorphisms p_z allow us to define the action of $U_q(\widehat{\mathfrak{sl}_2})$ in V_μ . The obtained $\mu + 1$ -dimensional representation of $U_q(\widehat{\mathfrak{sl}_2})$ is called the evaluation representation and is denoted by $V_\mu(z)$.

The Verma module $V_{\lambda,k}$ can be made a $U_q(\widehat{\mathfrak{sl}_2})$ -module by setting $Dv = v$, where $v \in V_{\lambda,k}$ is the highest weight vector. This condition uniquely determines the action of $D^{\pm 1}$ in $V_{\lambda,k}$. Let us say that a vector $w \in V_{\lambda,k}$ is at level n if $Dw = q^n w$. Denote the space of all such vectors by $V_{\lambda,k}[n]$. The subspace of top level vectors $V_{\lambda,k}[0]$ is a $U_q(\mathfrak{sl}_2)$ -subrepresentation isomorphic to the Verma module $\mathcal{M}_\lambda$ with highest weight λ over $U_q(\mathfrak{sl}_2)$.

We are interested in formal expressions of the form

$$(3.10) \quad \Phi(z) = \sum_{n \in \mathbb{Z}} \Phi[n] z^{-n},$$

where $\Phi[m] : V_{\lambda,k}[n] \rightarrow V_{\nu,k}[n+m] \otimes V_\mu(z)$ are linear maps such that

$$(3.11) \quad \Phi(z)au = \Delta(a)\Phi(z)u, \quad a \in U_q(\widehat{\mathfrak{sl}_2}), u \in V_{\lambda,k}.$$

Affording a slight abuse of terminology, we can say that $\Phi(z)$ is a $U_q(\widehat{\mathfrak{sl}_2})$ intertwining operator $V_{\lambda,k} \rightarrow V_{\nu,k} \otimes V_\mu(z)$.³ It is easy to show that any such operator is uniquely determined by its action on top level vectors. Moreover, it is clear that if $w \in V_{\lambda,k}$ is top level then it is enough to know the top level component of $\Phi(z)w$ in order to retrieve $\Phi(z)w$. This shows that $\Phi(z)$ is uniquely determined by the map $\Phi[0] : V_{\lambda,k}[0] \rightarrow V_{\nu,k}[0] \otimes V_\mu(z)$. This map must be an intertwining operator $\mathcal{M}_\lambda \rightarrow \mathcal{M}_\nu \otimes V_\mu$ over $U_q(\mathfrak{sl}_2)$. Moreover, for generic values of q, k this condition is necessary and sufficient for $\Phi[0]$ to extend to the entire module $V_{\lambda,k}$ [FR]. From now on we consider this generic situation.

Clebsch-Gordan formula tells us that there is a unique nonzero intertwiner $\Phi^{\lambda,\nu,\mu}$ of form (3.10) up to a constant if $\nu + \mu - \lambda$ is even, nonnegative, and not greater than 2μ , and no nonzero intertwiners of this form otherwise.

Let $u \in V_\mu(z)^*$. Define the operator $\Phi^{\lambda,\nu,\mu}(u, z) : V_{\lambda,k} \rightarrow \hat{V}_{\nu,k}$, $\Phi^{\lambda,\nu,\mu}(u, z)w = u(\Phi^{\lambda,\nu,\mu}(z)w)$.

Despite the range of the operator $\Phi^{\lambda,\nu,\mu}(u, z)$ lies in the completion of the highest weight module $V_{\nu,k}$, one can form products of such operators. If $|z_1| > |z_2| > \dots > |z_N|$ then the product

$$(3.12) \quad \Phi^{\lambda_1, \lambda_0, \mu_1}(u_1, z_1) \dots \Phi^{\lambda_N, \lambda_{N-1}, \mu_N}(u_N, z_N)$$

is a well defined linear map: $V_{\lambda_N, k} \rightarrow \hat{V}_{\lambda_0, k}$.

Let $v_{\lambda_N, k}$ and $v_{\lambda_0, k}^*$ be the highest weight vector of $V_{\lambda_N, k}$ and the lowest weight vector of $V_{\lambda_0, k}^*$, respectively, and let $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_N)$. Form the scalar product

$$(3.13) \quad \varphi^\Lambda(u_1, \dots, u_N, z_1, \dots, z_N) = \langle v_{\lambda_0, k}^*, \Phi^{\lambda_1, \lambda_0, \mu_1}(u_1, z_1) \dots \Phi^{\lambda_N, \lambda_{N-1}, \mu_N}(u_N, z_N) v_{\lambda_N, k} \rangle.$$

³This statement is not quite precise because if $u \in V_{\lambda,k}$ and $z \in \mathbb{C}^*$ then $\Phi(z)u$ lies in $\hat{V}_{\lambda,k} \otimes V_\mu(z)$, where $\hat{V}_{\lambda,k}$ is the completion of $V_{\lambda,k}$ which allows infinite sums of homogeneous

This scalar product is a matrix element of the intertwiner (3.12). We will regard it as a holomorphic function in $z_1, \dots, z_N$ in the region $|z_1| > |z_2| > \dots > |z_N|$ with values in the finite dimensional space $V_{\mu_1} \otimes \dots \otimes V_{\mu_N}$ and write it as $\varphi^\Lambda(z_1, \dots, z_N)$.

Let $h(\lambda) = \frac{\lambda^2 + 2\lambda}{2(k+2)}$. Let $\Delta_i(\Lambda) = h(\lambda_{i-1}) - h(\lambda_i)$, $1 \leq i \leq N$. Define a new (multivalued) function

$$(3.14) \quad \Psi^\Lambda(z_1, \dots, z_N) = z_1^{\Delta_1(\Lambda)} \dots z_N^{\Delta_N(\Lambda)} \varphi^\Lambda(z_1, \dots, z_N).$$

It turns out that the function Ψ^Λ is a product of a scalar function and a solution of the quantum KZ equations associated with a certain system of trigonometric R -matrices which is described explicitly as follows.

Let $m, n \geq 0$ be integers. As a $U_q(\mathfrak{sl}_2)$ -module, $V_m \otimes V_n$ decomposes as

$$V_m \otimes V_n = \bigoplus_{r=0}^{\min(m,n)} V_{m+n-2r}.$$

Let u_m be a highest weight vector of V_m , and let ω_{mn}^r be the highest weight vectors of the components V_{m+n-2r} such that $(e \otimes 1)\omega_{mn}^r = \omega_{mn}^{r-1}$ for $r \geq 1$, and $\omega_{mn}^0 = u_m \otimes u_n$. Let $P_{mn}^r : V_n \otimes V_m \rightarrow V_m \otimes V_n$ be the $U_q(\mathfrak{sl}_2)$ -invariant map such that $P_{mn}^r \omega_{nm}^r = \omega_{mn}^r$ and $P_{mn}^r \omega_{nm}^l = 0$ if $r \neq l$.

Define the map $R_{ij}^q(z) : V_{\mu_i} \otimes V_{\mu_j} \rightarrow V_{\mu_i} \otimes V_{\mu_j}$ by

$$(3.15) \quad R_{ij}^q(z) = \sum_{r=0}^{\min(\mu_i, \mu_j)} \prod_{l=0}^{r-1} \frac{1 - zq^{\mu_i + \mu_j - 2l}}{z - q^{\mu_i + \mu_j - 2l}} P_{\mu_i \mu_j}^r \sigma_{\mu_i \mu_j},$$

where $\sigma_{\mu_i \mu_j} : V_{\mu_i} \otimes V_{\mu_j} \rightarrow V_{\mu_j} \otimes V_{\mu_i}$ is the permutation of factors.

Let $W = V_{\mu_1} \otimes \dots \otimes V_{\mu_N}$. Fix a complex number λ_0 . Let $v_i \in V_{\mu_i}$, $1 \leq i \leq N$, and let $Kv_i = q^{m_i} v_i$. Set $m = \sum_{j=1}^N m_j$. Define the operators $A_i^{q, \lambda_0} : W \rightarrow W$ by

$$(3.16) \quad A_i^{q, \lambda_0}(v_1 \otimes v_2 \otimes \dots \otimes v_N) = q^{(2\lambda_0 + 2 + m)m_i} v_1 \otimes v_2 \otimes \dots \otimes v_N.$$

Proposition 3.1. [CP]

(i) $\{R_{ij}^q(z)\}$ is a system of trigonometric R -matrices on $V_{\mu_1}, \dots, V_{\mu_N}$, regular if $|q| < 1$.

(ii) $\{A_i^{q, \lambda_0}\}$ are compatible to $\{R_{ij}^q(z)\}$.

Thus, the matrices $\{R_{ij}^q(z)\}$ and $\{A_i^{q, \lambda_0}\}$ define a holomorphic vector bundle with fiber W . We will denote this bundle by E_N^{q, λ_0} .

Observe that $E_N^{q, \lambda_0} = \bigoplus_{r=0}^{\mu} E_{N,r}^{q, \lambda_0}$, where $E_{N,r}^{q, \lambda_0}$ is the subbundle of E_N^{q, λ_0} whose fiber is the subspace W_r of vectors of weight $\mu - 2r$ in W , $\mu = \sum_{j=1}^N \mu_j$.

Because the transition matrices of the bundle $E_{N,r}^{q, \lambda_0}$ depend only on the ratios z_i/z_j , this bundle can be obtained from a bundle over T^{N-1} . Indeed, let $\Theta \subset T^N$ be the diagonal: $\Theta = \{(z, z, \dots, z) | z \in T\}$, and let η be the projection: $\eta : T^N \rightarrow T^N/\Theta$ (the space T^N/Θ is isomorphic to T^{N-1}). Denote by ξ the map $T^N \rightarrow T$ which acts according to the formula $\xi(z_1, \dots, z_N) = z_1 z_2 \dots z_N$. Let $\Delta_0 = \frac{(2\lambda_0 + 2 + r)r}{2N(k+2)}$.

Then $E_{N,r}^{q, \lambda_0} = \xi^*(L_{\Delta_0}) \otimes \eta^*(\hat{E}_{N,r}^{q, \lambda_0})$, where $\hat{E}_{N,r}^{q, \lambda_0}$ is a holomorphic vector bundle on T^{N-1}/Θ .

Proposition 3.2. [FR] *The function $\Psi^\Lambda(z_1, \dots, z_N)$ can be represented in the form*

$$(3.17) \quad \Psi^\Lambda(z_1, \dots, z_N) = \prod_{i < j} G_{\mu_i \mu_j} \left(\frac{z_i}{z_j} \right) \psi^\Lambda(z_1, \dots, z_N),$$

where ψ^Λ is a solution of the quantum KZ equations (2.5) regular in the region $|z_1| \geq |z_2| \geq \dots \geq |z_N|$, and $G_{\mu_i \mu_j}(z)$ is a scalar-valued meromorphic function in $\mathbb{C}^*$.

The functions $G_{\mu_i \mu_j}(z)$ are described in [FR] as certain infinite products. In this paper, they will not be of further interest.

Now let $s = t_{j_1} \dots t_{j_n}$ be a permutation. Let $s_l = t_{j_1} \dots t_{j_l}$, $1 \leq l \leq n$. Define the function

$$(3.18) \quad \begin{aligned} \varphi^{s, \Lambda}(u_1, \dots, u_N, z_1, \dots, z_N) &= R_{s_{n-1}(j_n) s_{n-1}(j_n+1)}^q \dots R_{s_1(j_2) s_1(j_2+1)}^q R_{j_1, j_1+1}^q \times \\ &< v_{\lambda_0, k}^*, \Phi^{\lambda_1, \lambda_0, \mu_{s(1)}}(u_{s(1)}, z_{s(1)}) \dots \Phi^{\lambda_N, \lambda_{N-1}, \mu_{s(N)}}(u_{s(N)}, z_{s(N)}) v_{\lambda_N, k} >. \end{aligned}$$

(Abusing notation, we write R_{ij} instead of $R_{ij}(z_i/z_j)$). Define $\Psi^{s, \Lambda}$ by

$$(3.19) \quad \Psi^{s, \Lambda}(z_1, \dots, z_N) = z_{s(1)}^{\Delta_1(\Lambda)} \dots z_{s(N)}^{\Delta_N(\Lambda)} \varphi^{s, \Lambda}(z_1, \dots, z_N)$$

and $\psi^{s, \Lambda}$ by

$$(3.20) \quad \Psi^{s, \Lambda}(z_1, \dots, z_N) = \prod_{s^{-1}(i) < s^{-1}(j)} G_{\mu_i \mu_j} \left(\frac{z_i}{z_j} \right) \psi^{s, \Lambda}(z_1, \dots, z_N),$$

Proposition 3.3. [FR] *$\psi^{s, \Lambda}$ satisfies the quantum KZ equations and is regular in the region $|z_{s(1)}| \geq |z_{s(2)}| \geq \dots \geq |z_{s(N)}|$.*

Assume $|q| < 1$. Using Proposition 3.1 and the results of Section 2, we deduce

Proposition 3.4.

- (i) $\tilde{\psi}^{s, \Lambda}$ is a holomorphic section of the bundle $E_N^{q, \lambda_0}(s^{-1}\Delta(\Lambda))$.
- (ii) For any $\lambda \in \mathbb{C}$, there exist exactly $d = \dim W$ vectors Λ such that $\lambda_0 = \lambda$ and $\varphi^\Lambda \neq 0$. These vectors can be arranged in an order $\Lambda_1(s), \dots, \Lambda_d(s)$ so that $\tilde{\psi}^{s, \Lambda_i(s)} = \tilde{\psi}^{s, i}$, $1 \leq i \leq d$, where $\tilde{\psi}^{s, i}$, $s \in S_N$, is defined by (2.13).

Thus, we have shown that matrix elements of intertwining operators between representations of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ can be geometrically interpreted as holomorphic sections of a certain holomorphic vector bundle. This interpretation remains valid if $U_q(\widehat{\mathfrak{sl}}_2)$ is replaced with $U_q(\widehat{\mathfrak{g}})$ where $\mathfrak{g}$ is an arbitrary simple Lie algebra.

4. Connection matrices as clutching transformations.

Let $s_1, s_2 \in S_N$. Then the systems of sections $\tilde{\psi}^{(s_1, i_1)}$ and $\tilde{\psi}^{(s_2, i_2)}$ are related by a connection matrix $C^{s_1 s_2} = \{c_{i_1 i_2}^{s_1 s_2}\}$, where $c_{i_1 i_2}^{s_1 s_2}$ is a meromorphic section of the bundle $\mathcal{B}(\Delta^{(s_1, i_1)} - \Delta^{(s_2, i_2)})$:

$$(4.1) \quad \tilde{\psi}^{s_1, i_1} = \sum_{i_2}^d c_{i_1 i_2}^{s_1 s_2} \otimes \tilde{\psi}^{s_2, i_2}.$$

The connection matrices will have poles since the systems of solutions $\psi^{(s,i)}$, $1 \leq i \leq d$, are not everywhere linear independent. According to Proposition 2.5, the poles will be at points where $\frac{z_{s(j_1)}}{z_{s(j_2)}}p^n$ is a pole of $R_{s(j_1)s(j_2)}(z)$ for a suitable $n \in \mathbb{Z}$ and pair of indices j_1, j_2 such that $j_1 > j_2$.

Thus, any matrix element of the connection matrix can be written as a product of powers of $z_1, \dots, z_N$ and a rational expression of elliptic functions in $\log z_1, \dots, \log z_N$.

Let $s \in S_N$ and $s' = s \cdot (ij)$, where (ij) is the transposition of i and j . Let R_{ij}^q and A_i be defined by (3.15) and (3.16).

Proposition 4.1. [FR] *There exists a system of meromorphic functions $B_{ij}(\zeta)$ with values in $\text{Mat}_d(\mathbb{C})$, $1 \leq i, j \leq N$ such that $C^{s,s'}(z_1, \dots, z_N) = B_{s(i)s(j)}(\log \frac{z_{s(i)}}{z_{s(j)}})$. These matrices satisfy the conditions:*

(i) *the quantum Yang-Baxter equation:*

$$(4.2) \quad B_{ij}(\zeta_1 - \zeta_2) B_{ik}(\zeta_1 - \zeta_3) B_{jk}(\zeta_2 - \zeta_3) = B_{jk}(\zeta_2 - \zeta_3) B_{ik}(\zeta_1 - \zeta_3) B_{ij}(\zeta_1 - \zeta_2);$$

(ii) *unitarity:*

$$(4.3) \quad B_{ij}(\zeta) = B_{ji}^{-1}(-\zeta);$$

(iii) *double periodicity:*

$$(4.4) \quad B_{ij}(\zeta + \log p) = B_{ij}(\zeta); \quad B_{ij}(\zeta + 2\pi\sqrt{-1}) = LB_{ij}(\zeta)L',$$

where $L, L' \in \text{End}(W)$ are constant diagonal matrices.

This statement shows that the connection matrices provide elliptic solutions to the quantum Yang-Baxter equation (elliptic quantum R -matrices).

Now we can give an alternative construction of the bundle E_N^{q,λ_0} , using the connection matrices as clutching transformations.

Let X be a complex analytic space, and let $\{U_i, i \in I\}$ be an open cover of X . Let $\mathcal{E}_i \rightarrow U_i$ be holomorphic vector bundles, and let $\beta_{ij} : \mathcal{E}_i|_{U_i \cap U_j} \rightarrow \mathcal{E}_j|_{U_i \cap U_j}$ be isomorphisms of holomorphic bundles such that the consistency conditions $\beta_{ij}\beta_{ji} = \text{id}$ in $U_i \cap U_j$, $\beta_{ij}\beta_{jk}\beta_{ki} = \text{id}$ in $U_i \cap U_j \cap U_k$ are satisfied. Then one can construct a holomorphic vector bundle $\mathcal{E}$ on X by setting $\mathcal{E}|_{U_i} = \mathcal{E}_i|_{U_i}$ and defining the clutching transformation from U_i to U_j to be β_{ij} .

Assume that $p^n/q^m \neq 1$ for any nonzero integers n and m . Let $s \in S_N$, and let H_s be the set of all points $(z_1, \dots, z_N) \in T^N$ such that if $n \in \mathbb{Z}$ and $j_1 > j_2$ then $\frac{z_{s(j_1)}}{z_{s(j_2)}}p^n$ is not a pole of $R_{s(j_1)s(j_2)}^q(z)$. The sections $\{\tilde{\psi}^{s,i}, 1 \leq i \leq d\}$ are linearly independent over H_s .

Lemma 4.2. $\{H_s, s \in S_N\}$ is an open cover of T^N .

Proof. Let $P = (z_1, \dots, z_N) \in T^N$. We need to show that there exists $s \in S_N$ such that $P \in H_s$. Let Q be the infinite cyclic subgroup in T generated multiplicatively by q . Let $x_1, \dots, x_r \in T/Q$ be the distinct images of the points $z_1, \dots, z_N \in T$ under the homomorphism $h : T \rightarrow T/Q$, and let $X_j = h^{-1}(x_j)$, $1 \leq j \leq r$. Inside X_j , the elements are naturally ordered: for $a, b \in X_j$, we say that $a \preccurlyeq b$ if and only if $a/b = q^m p^n$, where $m, n \in \mathbb{Z}$ and $m > 0$. Let $s \in S_N$ be a permutation such that $z_{s(i)} \preccurlyeq z_{s(j)}$ implies $i \leq j$. Then $P \in H_s$. ■

Now define a holomorphic vector bundle $\mathcal{E}_N^{q,\lambda_0}$ on T^N as follows. Set $\mathcal{E}_N^{q,\lambda_0}|_{H_s}$ to be isomorphic to $\bigoplus_{i=1}^d \mathcal{B}(-\Delta^{(s,i)})|_{H_s}$, and define the clutching transformations

$$\beta_{s_1 s_2} : \bigoplus_{i=1}^d \mathcal{B}(-\Delta^{(s_1,i)})|_{H_{s_1} \cap H_{s_2}} \rightarrow \bigoplus_{i=1}^d \mathcal{B}(-\Delta^{(s_2,i)})|_{H_{s_1} \cap H_{s_2}}$$

by $\beta_{s_1 s_2} = C^{s_2 s_1}$. It is trivial to check that the consistency conditions are satisfied

Proposition 4.3. *The bundles E_N^{q,λ_0} and $\mathcal{E}_N^{q,\lambda_0}$ are isomorphic to each other.*

Proof. The system of sections $\{\tilde{\psi}^{s,i}, 1 \leq i \leq d\}$ defines an isomorphism between the bundles $\oplus_{i=1}^d \mathcal{B}(-\Delta^{(s,i)})|_{H_s}$ and $E_N^{q,\lambda_0}|_{H_s}$. Transition from one system of sections to another is performed by the connection matrix $C^{s_1 s_2}$. ■

5. Topology of the quantum Knizhnik-Zamolodchikov equations and crystal bases.

In this section we study the vector bundle E_N^{q,λ_0} topologically, disregarding its holomorphic structure. It turns out that the topological structure of E_N^{q,λ_0} is quite nontrivial and can be described in terms of the combinatorics of crystal bases in representations of quantum groups.

Clearly, we may assume, without loss of generality, that $\dim V_{\mu_j} \geq 2$ for all j . Then $\text{rank}(E_N^{q,\lambda_0}) \geq 2^N$.

Recall some properties of characteristic classes. Let $\mathcal{E}$ be a vector bundle on T^N . Let $c_k(\mathcal{E}) \in H^{2k}(T^N, \mathbb{Z})$, $0 \leq k \leq N$, be the Chern classes of $\mathcal{E}$ (by convention $c_0 = 1$). The sum $c(\mathcal{E}) = \sum_{k=0}^N c_k(\mathcal{E}) \in \oplus_{k=0}^N H^{2k}(T^N, \mathbb{Z})$ is called the total Chern class of $\mathcal{E}$. It has the property $c(\mathcal{E} \oplus \mathcal{F}) = c(\mathcal{E})c(\mathcal{F})$ for any two vector bundles $\mathcal{E}$ and $\mathcal{F}$. Also, if $\mathcal{E}$ and $\mathcal{F}$ are one-dimensional then $c_1(\mathcal{E} \otimes \mathcal{F}) = c_1(\mathcal{E}) + c_1(\mathcal{F})$ [Mi].

Let $f_j(x_1, \dots, x_N)$ be the elementary symmetric polynomials:

$$\sum_{j=0}^N f_j t^{N-j} = \prod_{j=1}^N (t + x_j).$$

Let $Q_k(y_1, \dots, y_k)$ be the Newton polynomials defined by $Q_k(f_1, \dots, f_k) = \sum_{j=1}^N x_j^k$. The Chern character of a vector bundle $\mathcal{E}$ is defined by

$$(5.1) \quad \text{Ch}(\mathcal{E}) = r + \sum_{k=1}^N \frac{1}{k!} Q_k(c_1, \dots, c_k),$$

where r is the rank of $\mathcal{E}$. It has the properties $\text{Ch}(\mathcal{E} \oplus \mathcal{F}) = \text{Ch}(\mathcal{E}) + \text{Ch}(\mathcal{F})$, $\text{Ch}(\mathcal{E} \otimes \mathcal{F}) = \text{Ch}(\mathcal{E})\text{Ch}(\mathcal{F})$.

Proposition 5.1. *A complex vector bundle over T^N of rank $r \geq N$ is uniquely determined by its Chern classes.*

Proof. Two bundles of rank $r \geq N$ over T^N are equivalent if and only if they are stable equivalent. Therefore, a bundle of rank $\geq N$ is uniquely determined by its class in the ring $K(T^N)$. A well known theorem of K -theory ([K, Theorem 5.3.25]), asserts that the Chern character induces an isomorphism of rings: $K(T^N) \otimes \mathbb{Q} \rightarrow H^{2*}(T^N) \otimes \mathbb{Q}$. The complex K -ring of the torus is isomorphic to the even part of its cohomology ring. This fact follows from the rule of evaluation of the K -ring of a product of two spaces (see Prop 4.3.24 in [K]). Therefore, $K(T^N)$ is a torsion free abelian group, which implies that the natural map $K(T^N) \rightarrow K(T^N) \otimes \mathbb{Q}$ is a monomorphism. Thus, the composed map $\text{Ch} : K(T^N) \rightarrow H^{2*}(T^N, \mathbb{Q})$ is injective, Q.E.D. ■

Since $\text{rank}(E_N^{q,\lambda_0}) > N$, in order to describe the bundle E_N^{q,λ_0} topologically it is enough to calculate $c(E_N^{q,\lambda_0})$. This turns out to be quite simple.

When q goes to 0, the operators P_{mn}^r defined in Chapter 3 tend to finite limits – a remarkable phenomenon known as crystallization [Ka]. This implies that the B matrices $B^q(x)$ have finite limits as $q \rightarrow 0$.

It is clear that the topological structure of the bundle $E_{N,r}^{q,\lambda_0}$ is stable under deformations and hence independent of the values of q and λ_0 (as long as $|q| < 1$). Therefore, E_N^{q,λ_0} is topologically equivalent to the bundle $\tilde{E}_N^q = \bigoplus_{r=0}^{\mu} E_{N,r}^{q,-1-r/2}$. As $q \rightarrow 0$, the transition matrices of the bundle $\tilde{E}_N^q$ have finite limits: $R_{ij}^q \rightarrow R_{ij}^0$, $A_i^{q,-1-r/2} \rightarrow 1$ on W_r . Therefore, there exists a limiting bundle $\tilde{E}_N^0$, with the fiber W , defined by the matrices $R_{ij}(z) = R_{ij}^0(z)$ and $A_i = 1$ according to (1.4) and (1.5), and the bundle E_N^{q,λ_0} is topologically equivalent to $\tilde{E}_N^0$ for any q and λ_0 .

Let us now describe the limiting R -matrix R_{ij}^0 . Let $u_m^l = f^l u_m \in V_m$, $0 \leq l \leq m$.

Proposition 5.2. [Ka] *Let $q = 0$. Then $P_{mn}^r \sigma(u_m^i \otimes u_n^j) = 0$ unless $\min(i, n-j) = r$. If $\min(i, m-j) = r$ then*

$$(5.2) \quad P_{mn}^r \sigma(u_m^i \otimes u_n^j) = \begin{cases} u_m^j \otimes u_n^i, & i+j \leq m \\ u_m^{m-i} \otimes u_n^{2i+j-m}, & m < i+j \leq n \\ u_m^{n-m+j} \otimes u_n^{m-n+i}, & n < i+j \leq n+m \end{cases}$$

if $m \leq n$, and

$$(5.3) \quad P_{mn}^r \sigma(u_m^i \otimes u_n^j) = \begin{cases} u_m^j \otimes u_n^i, & i+j \leq n \\ u_m^{i+2j-n} \otimes u_n^{n-j}, & m < i+j \leq n \\ u_m^{n-m+j} \otimes u_n^{m-n+i}, & m < i+j \leq n+m \end{cases}$$

if $m > n$.

Corollary 5.3. *The R -matrix at $q = 0$ has the form*

$$(5.4) \quad R_{ij}^0(z)(u_{\mu_i}^k \otimes u_{\mu_j}^l) = \begin{cases} z^{-\min(k, \mu_j-l)} u_{\mu_i}^l \otimes u_{\mu_j}^k, & k+l \leq \mu_i \\ z^{-\min(k, \mu_j-l)} u_{\mu_i}^{\mu_i-k} \otimes u_{\mu_j}^{2k+l-\mu_i}, & \mu_i < k+l \leq \mu_j \\ z^{-\min(k, \mu_j-l)} u_{\mu_i}^{\mu_j-\mu_i+l} \otimes u_{\mu_j}^{\mu_i-\mu_j+k}, & \mu_j < k+l \leq \mu_i + \mu_j \end{cases}$$

if $\mu_i \leq \mu_j$, and

$$(5.5) \quad R_{ij}^0(z)(u_{\mu_i}^k \otimes u_{\mu_j}^l) = \begin{cases} z^{-\min(k, \mu_j-l)} u_{\mu_i}^l \otimes u_{\mu_j}^k, & k+l \leq \mu_j \\ z^{-\min(k, \mu_j-l)} u_{\mu_i}^{k+2l-\mu_j} \otimes u_{\mu_j}^{\mu_j-l}, & \mu_j < k+l \leq \mu_i \\ z^{-\min(k, \mu_j-l)} u_{\mu_i}^{\mu_j-\mu_i+l} \otimes u_{\mu_j}^{\mu_i-\mu_j+k}, & \mu_i < k+l \leq \mu_i + \mu_j \end{cases}$$

if $\mu_i > \mu_j$.

Let $L = (l_1, \dots, l_N)$, $0 \leq l_j \leq \mu_j$. Let $U_L = u_{\mu_1}^{l_1} \otimes \dots \otimes u_{\mu_N}^{l_N} \in W$. The vectors $\{U_L\}$ form a basis of W (the crystal basis).

Corollary 5.4. *In the basis $\{U_L\}$, the operator $M_j(z_1, \dots, z_N)$ defined by (2.7) can be represented in the form of a product*

$$(5.6) \quad M_j(z_1, \dots, z_N) = Y_j(z_1, \dots, z_N) \pi_j$$

where π_j is a permutation matrix ($\pi_j \in S_d$, $d = \dim W$), and Y_j is a diagonal matrix whose eigenvalues are products of integer powers of $z_1, \dots, z_N$.

It is clear that the permutations π_j commute with each other. Let α_j be the order of π_j in S_d . Let $\hat{\Gamma}^N$ be the torus $\prod_{j=1}^N (\mathbb{C}^* / \Gamma_j)$, where Γ_j is the infinite cyclic

group multiplicatively generated by p^{α_j} . Let $\beta : \hat{T}^N \rightarrow T^N$ be the natural covering. Consider the holomorphic vector bundle $\beta^*(\tilde{E}_N^0)$ on $\hat{T}^N$. Corollary 5.4 implies that this bundle breaks up into a direct sum of line bundles, and each of these line bundles is a tensor product of line bundles over factors $\mathbb{C}^*/\Gamma_j$. This allows us to calculate $c(\beta^*(\tilde{E}_N^0))$ using the properties of Chern classes that we discussed above. Applying the map $(\beta^*)^{-1} : H^*(\hat{T}^N, \mathbb{Q}) \rightarrow H^*(T^N, \mathbb{Q})$ to $c(\beta^*(\tilde{E}_N^0))$, we obtain the sought-for total Chern class $c(E_N^0)$.

For the sake of brevity, we implement this calculation under the assumption that all V_{μ_j} are the same: $\mu_j = m$ for all j . Then formulas (5.4) and (5.5) undergo a major simplification: $R_{ij}^0(z) = R^0(z)$, $i \neq j$, and

$$(5.7) \quad R^0(z)(u_m^k \otimes u_m^l) = z^{-\min(k, m-l)} u_m^l \otimes u_m^k.$$

In the formulas below, indices i, j are allowed to take values between 1 and N , and operations on them are fulfilled modulo N . For instance, $N+1$ is identified with 1, and 0 is identified with N .

Introduce the notation $I(k, l) = \min(k, m-l)$, $J(L, j) = \sum_{i=1}^{j-1} I(l_i, l_{i+1})$, and $J(L) = \sum_{i=1}^N I(l_i, l_{i+1})$.

Substituting (5.7) into (2.6), we obtain

$$(5.8) \quad M_j(z_1, \dots, z_N) u_m^{l_1} \otimes \dots \otimes u_m^{l_{N-1}} \otimes u_m^{l_N} = z_j^{-J(L) + I(l_{j-1}, l_j)} \prod_{i \neq j} z_i^{I(l_{i-1}, l_i)} p^{-J(L, j)} u_m^{l_N} \otimes u_m^{l_1} \otimes \dots \otimes u_m^{l_{N-1}}$$

This shows that for all j π_j is the cyclic permutation $(123\dots N)$. Therefore, $\alpha_j = N$ for every j . Let

$$(5.9) \quad \hat{M}_j(z_1, \dots, z_N) = M_j(z_1, \dots, p^{N-1} z_j, \dots, z_N) \dots M_j(z_1, \dots, p z_j, \dots, z_N) M_j(z_1, \dots, z_j, \dots, z_N)$$

The matrices $\hat{M}_j$ are the transition matrices for the bundle $\beta^*(\tilde{E}_N^0)$. On the other hand, as we have mentioned, these matrices are diagonal:

$$(5.10) \quad \hat{M}_j(z_1, \dots, z_N) U_L = z_j^{-(N-1)J(L)} \left(\prod_{i \neq j} z_i \right)^{J(L)} p^{K(L)} U_L,$$

where $K(L)$ is an integer.

Let $x_i = \frac{1}{\log|p|} \operatorname{Re} \log z_i$, $y_i = \frac{1}{2\pi\sqrt{-1}} \operatorname{Im} \log z_i$. Let $[\Omega] \in H^2(T^N, \mathbb{Z})$ be the cohomology class of the differential form

$$(5.11) \quad \Omega = N \sum_{j=1}^N dx_j \wedge dy_j - \left(\sum_{j=1}^N dx_j \right) \wedge \left(\sum_{j=1}^N dy_j \right)$$

Formula (5.10) immediately yields an expression for $c(\beta^*(\tilde{E}_N^0))$:

$$(5.12) \quad c(\beta^*(\tilde{E}_N^0)) = \prod_L (1 + J(L)[\Omega]),$$

where the product is taken over all $L = (l_1, \dots, l_N)$ with $1 \leq l_j \leq m$. This, in turn, gives a formula for $c(\tilde{E}_N^0)$:

$$(5.13) \quad c(\tilde{E}_N^0) = \prod_L \left(1 + \frac{J(L)}{N} [\Omega] \right).$$

Remark 5.1. Formula (5.12) contains fractions, but it always gives an integral cocycle – the total Chern class of a vector bundle.

Thus, we have proved

Proposition 5.5. *If $V_{\mu_i} = V_m$ for $1 \leq i \leq N$ then*

$$(5.14) \quad c(E_{N,r}^{q,\lambda_0}) = \prod_{L: \sum l_j = r} (1 + \frac{J(L)}{N}[\Omega]), \quad 0 \leq r \leq Nm,$$

$$(5.15) \quad c(E_N^{q,\lambda_0}) = \prod_L (1 + \frac{J(L)}{N}[\Omega]),$$

for all values of q and λ_0 .

6. Special case $N = 2$.

According to Chapter 3, in the special case $N = 2$ the bundle $E_{N,r}^{q,\lambda_0}(\Delta)$ can be represented as a product:

$$(6.1) \quad E_{2,r}^{q,\lambda_0}(\Delta) = \xi^*(L_{\Delta_0 + \frac{1}{2}(\Delta_1 + \Delta_2)}) \otimes \eta^*(\hat{E}_{2,r}^{q,\lambda_0}(\frac{\Delta_1 - \Delta_2}{2})),$$

where $\xi, \eta : T^2 \rightarrow T$ are defined by $\xi(z_1, z_2) = z_1 z_2$, $\eta(z_1, z_2) = z_1 / z_2$, $\Delta_0 = \frac{(2\lambda_0 + 2 + r)r}{4(k+2)}$, and $\hat{E}_{2,r}^{q,\lambda_0}$ is a bundle over T . Clearly, it is enough to understand $\hat{E}_{2,r}^{q,\lambda_0}$ in order to understand $E_{2,r}^{q,\lambda_0}$.

Proposition 6.1.

$$(6.2) \quad \begin{aligned} \text{rank}(\hat{E}_{2,r}^{q,\lambda_0}) = \rho_0 &= \begin{cases} r + 1, & r \leq \min(\mu_1, \mu_2) \\ \min(\mu_1, \mu_2) + 1, & \min(\mu_1, \mu_2) < r \leq \max(\mu_1, \mu_2) \\ \mu_1 + \mu_2 - r + 1, & \max(\mu_1, \mu_2) < r \leq \mu_1 + \mu_2 \end{cases} \\ \text{deg}(\hat{E}_{2,r}^{q,\lambda_0}) = \delta_0 &= \frac{1}{2}\rho_0(\rho_0 - 1) \end{aligned}$$

This proposition follows directly from formulas (4.4) and (4.5). It completely describes the topological structure of the bundle $\hat{E}_{2,r}^{q,\lambda_0}$.

The bundle $\hat{E}_{2,r}^{q,\lambda_0}$ is a holomorphic vector bundle over an elliptic curve. Such bundles were completely classified by M. Atiyah in 1957 [A]. M. Atiyah showed that the set of indecomposable bundles of a fixed rank and degree over an elliptic curve T can be identified with T : if $\mathcal{E}$ is an indecomposable bundle then any other indecomposable bundle $\mathcal{F}$ of the same rank and degree can be represented in the form $\mathcal{F} = \mathcal{E} \otimes L$, where L is a line bundle of degree 0.

In 1962 D. Mumford introduced the notions of stability and semistability for vector bundles over curves [Mu;NS]. A vector bundle $\mathcal{E}$ over a curve X is called stable if for every proper nonzero subbundle $\mathcal{F} \subset \mathcal{E}$ $\frac{\text{deg}(\mathcal{F})}{\text{rank}(\mathcal{F})} < \frac{\text{deg}(\mathcal{E})}{\text{rank}(\mathcal{E})}$, and semistable if $\frac{\text{deg}(\mathcal{F})}{\text{rank}(\mathcal{F})} \leq \frac{\text{deg}(\mathcal{E})}{\text{rank}(\mathcal{E})}$. If $\mathcal{E}(t)$ is a holomorphic family of vector bundles on X parametrized by an analytic variety $\mathcal{T}$ then the sets $\{t \in \mathcal{T} | \mathcal{E}(t) \text{ is stable}\}$ and $\{t \in \mathcal{T} | \mathcal{E}(t) \text{ is semistable}\}$ are Zariski open in $\mathcal{T}$. However, if X is an elliptic curve, stable bundles of rank ρ and degree δ over X exist if and only if ρ and δ are relatively prime. On the contrary, semistable bundles over X exist for any values of ρ and δ . Thus, a generic vector bundle over an elliptic curve is semistable.

Here are some properties of stable and semistable bundles.

Lemma 6.2.

- (i) If ρ and δ are relatively prime, every semistable bundle of rank ρ and degree δ is stable.
- (ii) If $\mathcal{E}$ is a stable (semistable) bundle and L is a line bundle then $\mathcal{E} \otimes L$ is stable (semistable).
- (iii) If $L_1, \dots, L_n$ are line bundles of the same degree then the bundle $\mathcal{E} = L_1 \oplus \dots \oplus L_n$ is semistable.
- (iv) Let $\mathcal{E}$ be a vector bundle over a curve X , and let $\gamma : \hat{X} \rightarrow X$ be a finite covering. If $\gamma^*\mathcal{E}$ is a stable (semistable) vector bundle over $\hat{X}$ then $\mathcal{E}$ is also stable (semistable).

Proof. Properties (i) and (ii) are obvious.

(iii) Let $\deg(L_j) = \delta$. Let $\mathcal{F}$ be a subbundle of rank ρ in $\mathcal{E}$. Then $\Lambda^\rho \mathcal{F}$ is a line bundle embedded in $\Lambda^\rho \mathcal{E}$. Obviously, $\Lambda^\rho \mathcal{E} = \bigoplus_i B_i$, where B_i are line bundles of degree $\rho\delta$. This implies that for at least one j the projection $\Lambda^\rho \mathcal{F} \rightarrow B_j$ is not zero. This projection is a nonzero regular section of the bundle $\Lambda^\rho \mathcal{F}^* \otimes B_j$. Thus, $\deg(\Lambda^\rho \mathcal{F}^* \otimes B_j) \geq 0$, which implies that $\deg(\mathcal{F}) \leq \rho\delta$.

(iv) Pulling back by γ does not change the rank of the bundle and multiplies its degree by the number of sheets of the covering. Therefore, if $\mathcal{F} \subset \mathcal{E}$ violates the stability (semistability) condition for $\mathcal{E}$, so does $\gamma^*\mathcal{F}$ for $\gamma^*\mathcal{E}$. ■

Proposition 6.3. *The bundle $\hat{E}_{2,r}^{q,\lambda_0}$ is semistable for generic values of q and λ_0 .*

Remark 6.1. If $\hat{E}_{2,r}^{q,\lambda_0}$ is semistable then, according to Lemma 6.2, (ii), so is $\hat{E}_{2,r}^{q,\lambda_0}(\Delta)$ for every complex number Δ .

Proof. First let us show that for a generic value of q the bundle $\hat{E}_{2,r}^{q,-r-1/2}$ is semistable. Since the set of such values is automatically Zariski open in the unit disk, we only need to show that there exists at least one such value. We will prove it for $q = 0$.

Let $\hat{T} = \mathbb{C}^*/\Pi^2$, where Π^2 is the infinite cyclic subgroup multiplicatively generated by p^2 . Let $\beta : \hat{T} \rightarrow T$ be the natural 2-sheeted covering. Formulas (5.4) and (5.5) show that the bundle $\beta^* \hat{E}_{2,r}^{0,-r-1/2}$ is a direct sum of ρ_0 line bundles of degree $\rho_0 - 1$. By Lemma 6.2, (iii), this bundle is semistable. Then Lemma 6.2, (iv) ensures the semistability of $\hat{E}_{2,r}^{0,-r-1/2}$.

We know that the set of pairs (q, λ_0) for which the bundle $\hat{E}_{2,r}^{q,\lambda_0}$ is semistable is Zariski open in $\mathcal{D}_0 \times \mathbb{C}$, $\mathcal{D}_0$ being the open unit disk punctured at 0. This set is also nonempty because it contains almost every point of the form $(q, -r - 1/2)$. Hence, it contains almost every point of $\mathcal{D}_0 \times \mathbb{C}$. ■

Corollary 6.4. *If $r = 1$ then $\hat{E}_{2,r}^{q,\lambda_0}$ is stable, and hence indecomposable for generic values of q and λ_0 .*

Proof. The bundle $\hat{E}_{2,1}^{q,\lambda_0}$ has degree 1 and rank 2, and it is semistable for generic values of parameters. Hence, by Lemma 6.2, (i), it is (generically) stable and therefore indecomposable. ■

Proposition 6.5. *For generic q and λ_0 , the vector bundle $\hat{E}_{2,r}^{q,\lambda_0}$ is a direct sum of line bundles if ρ_0 is odd, and a sum of indecomposable 2-dimensional bundles if ρ_0 is even. Thus, $\hat{E}_{2,r}^{q,\lambda_0}$ is almost always a direct sum of stable bundles.*

Proof. Fix a complex number a and consider the family of vector bundles $\hat{E}_{2,r}^{q,-1-\frac{r}{2}+\frac{a}{\log q}}$, $|a| \leq 1$. This family is analytic in the unit disc, and at $a = 0$ the limiting bundle is

prescribed by (4) and (5) with $R_{12}(z) = R_{12}^0(z)$, $A_i(u_{\mu_1}^{m_1} \otimes u_{\mu_2}^{m_2}) = e^{2am_i} u_{\mu_1}^{m_1} \otimes u_{\mu_2}^{m_2}$. As before, $\beta^* \lim_{q \rightarrow 0} \hat{E}_{2,r}^{q, -1 - \frac{r}{2} + \frac{a}{\log q}}$ is a direct sum of line bundles. Moreover, it is easy to check that for a generic a , all these line bundles are pairwise non-isomorphic. This shows that such a decomposition takes place for $\beta^* \hat{E}_{2,r}^{q, \lambda_0}$ for generic values of q and λ_0 .

Now assume that $\hat{E}_{2,r}^{q, \lambda_0}$ is semistable but not a direct sum of stable bundles. Let $\hat{E}_{2,r}^{q, \lambda_0} = I_1 \oplus \cdots \oplus I_s$, where I_k are indecomposable. Suppose that I_k is unstable for some k . Then, according to [A], $I_k = \mathcal{E} \otimes \mathcal{F}$ where $\mathcal{E}$ is a stable bundle (of rank 1 if ρ_0 is odd and of rank 2 if ρ_0 is even), and $\mathcal{F}$ is an indecomposable degree zero bundle on T with $\text{rank}(\mathcal{F}) > 1$. Therefore, we have $\beta^* I_k = \beta^* \mathcal{E} \otimes \beta^* \mathcal{F}$. We know that $\beta^* \mathcal{E}$ is either a line bundle or a direct sum of two line bundles, whereas $\beta^* \mathcal{F}$ is obviously indecomposable. Thus, $\beta^* I_k$ is not a direct sum of line bundles. This does not happen for generic values of the parameters. ■

Remark 6.2. These statements indicate that the trigonometric R -matrices arising from $U_q(\widehat{\mathfrak{sl}_2})$, as a rule, give rise to 'generic' holomorphic bundles over T , i.e. ones corresponding to regular points of the moduli space of bundles. It would be interesting to check if this property holds for other simple Lie algebras and for $N > 2$.

Remark 6.3. It seems plausible that propositions 6.3 – 6.5 should hold for *arbitrary* values of q and λ_0 , not only for generic ones. However, a proof of this statement would probably have to involve a more delicate argument than what is used above.

Remark 6.4. It seems to be an interesting problem to separate the matrix elements of intertwining operators from the whole variety of holomorphic sections of the bundles $\hat{E}_{N,r}^{q, \lambda_0}(\Delta)$ by some geometric condition. It is not clear how to approach this question since, for instance, the bundle $\hat{E}_{2,r}^{q, \lambda_0}(\Delta)$, being generically semistable, has as many linearly independent regular sections as its degree, for all values of Δ .

7. Arbitrary root systems

In the recent paper [Ch] a quantum R -matrix R was interpreted as a 1-cocycle on the semidirect product $\tilde{W} = W \ltimes P$ of a Weyl group W of the type A_n and its weight lattice P with coefficients in a certain group $\mathcal{G}$ on which $\tilde{W}$ operates by automorphisms. The quantum KZ system can then be viewed as a condition on an element $g \in \mathcal{G}$ (regarded as a 0-cochain on $\tilde{W}$ with coefficients in $\mathcal{G}$) requiring that $dg(b) = R(b)$, $b \in P$, where d denotes the coboundary operator.

This interpretation is very useful since it can be made a definition of a quantum R -matrix and the KZ system if A_n is replaced with an arbitrary Dynkin diagram.

Our purpose now is to link the 1-cocycle and the vector bundle interpretations of the quantum KZ equations together and to extend the vector bundle interpretation to arbitrary root systems.

Let W be a (finite) Weyl group, and let P be its dual root lattice. Denote by $\mathcal{K}$ the field of P -periodic trigonometric functions on $P \otimes \mathbb{C}$ (i.e. rational expressions of monomials $e^{2\pi\sqrt{-1}\lambda(z)}$, $z \in P \otimes \mathbb{C}$, where $\lambda \in Q^\vee$, $Q^\vee$ being the dual lattice to P). Let $\mathcal{G} = G(\mathcal{K})$ where $G = GL(H)$ and H is a finite-dimensional representation of $\tilde{W} = W \ltimes P$.

Let $\tau \in \mathbb{C}^+$. We define the action of $\tilde{W}$ on $\mathcal{G}$ by $(wb \circ g)(z) = wg(w^{-1}z - b\tau)w^{-1}$, $b \in P$, $z \in P \otimes \mathbb{C}$, $w \in W$.

Let $S_0 \subset P \otimes \mathbb{R}$ be the set of points with a nontrivial stabilizer in $\tilde{W}$. Let $\mathcal{S} = P \otimes \mathbb{R} \setminus S_0 \subset P \otimes \mathbb{C}$. Let $\mathcal{G}_0$ be the subgroup of the elements of $\mathcal{G}$ that are

regular at S . Clearly, the action of $\tilde{W}$ maps elements of $\mathcal{G}_0$ to (other) elements of $\mathcal{G}_0$.

Definition 7.1. *A trigonometric quantum R -matrix on H is a W -invariant 1-cocycle R on $\tilde{W}$ with coefficients in $\mathcal{G}_0$ with the described action of $\tilde{W}$: for $x, y \in \tilde{W}$ $R(xy) = (y^{-1} \circ R(x))R(y)$ in $\mathcal{G}_0$.*

Trigonometric R -matrices defined in Chapter 1 (for $V_i = V$, $1 \leq i \leq N$) are obtained in the special case $H = V^{\otimes N}$, $W = S_N$, and the action of $\tilde{W}$ in H factorizes through the action of W in H by permutations of factors.

Let $\tilde{\mathcal{K}}$ be the field of meromorphic P -periodic functions in $P \otimes \mathbb{C}$, and let $\tilde{\mathcal{G}} = G(\tilde{\mathcal{K}})$. Let $\tilde{\mathcal{G}}_0$ be the subgroup of the elements of $\tilde{\mathcal{G}}$ that are regular at S . The action of $\tilde{W}$ in $\tilde{\mathcal{G}}_0$ is the same as in $\mathcal{G}_0$.

Definition 7.2. *Let $g \in \tilde{\mathcal{G}}_0$. The equations $(dg)(b) = R(b)$ with respect to g , for all $b \in P$, where d is the coboundary operator, are called the quantum KZ system for R .*

Again, the quantum KZ equations considered in Chapter 2 come about in the special case when $H = V^{\otimes N}$ and $W = S_N$.

Now, given a 1-cocycle R , we will construct a W -equivariant holomorphic vector bundle on the torus $T(W) = P \otimes \mathbb{C}/(P \oplus \tau P)$.

The set S partitions the space $P \otimes \mathbb{C}$ into infinitely many bounded chambers D_u labeled by elements $u \in \tilde{W}$, so that $u_2 \in \tilde{W}$ maps D_{u_1} to $D_{u_1 u_2}$. If two chambers D_{u_1} and D_{u_2} are adjacent (i.e the intersection of their boundaries has codimension 1) then $u_2 = u_1 s$ where $s \in \tilde{W}$ is a reflection. Consider the open cover of $P \otimes \mathbb{C}$ by the union of small neighborhoods of the chambers D_u . Let us define the transition matrix in the vicinity of the boundary between D_{u_1} and D_{u_2} (from D_{u_1} to D_{u_2}) to be

$$(7.1) \quad T_{u_1, s}(z) = (u_1^{-1} \circ R(u_1 s u_1^{-1}))(z), \quad z \in P \otimes \mathbb{C},$$

where the right hand side is regarded as a $GL(H)$ -valued function. By construction, this matrix is regular near the boundary. Because R is a 1-cocycle on $\tilde{W}$, the system of matrices $T_{u, s}$ is a holomorphic $GL(H)$ -valued Čech 1-cocycle on $P \otimes \mathbb{C}$. It is easy to check that this Čech cocycle is invariant under the action of $\tilde{W} \ltimes (P \oplus \tau P)$. Since every Čech cocycle automatically defines a vector bundle, we obtain a holomorphic vector bundle with fiber H on $P \otimes \mathbb{C}$, which descends to a holomorphic bundle over the torus $T(W) = P \otimes \mathbb{C}/(P \oplus \tau P)$.

In this setting, meromorphic solutions of the quantum KZ system are interpreted as meromorphic sections of the constructed vector bundle, since geometrically representation of a 1-cocycle as a coboundary of something is equivalent to constructing a global fundamental system of sections of the corresponding bundle.

In particular, in the special case $H = V^{\otimes N}$ and $W = S_N$, this scheme gives the vector bundle E_N constructed in Chapter 1 (assuming $A_1 A_2 \dots A_N = 1$) pushed forward to the $N - 1$ -dimensional torus T^N/Θ , where Θ is the main diagonal. This push-forward is possible since if $A_1 A_2 \dots A_N = 1$ then E_N is invariant under diagonal translations.

REFERENCES

- [Ao] Aomoto, K., *A note on holonomic q -difference system*, Algebraic analysis I, ed. by M.Kashiwara and T.Kawai, Acad Press, (1988), 25-28.
- [A] Atiyah, M., *Vector bundles over an elliptic curve*, Proc. Lond. Math. Soc. **7** (1957), 414-452.
- [CP] Chari, V., and Pressley, A., *Quantum affine algebras*, Comm. Math. Phys. **142** (1991), 261-

- [Ch] Cherednik, I.V., *Quantum Knizhnik-Zamolodchikov equations and affine root systems*, preprint (1992).
- [D] Drinfeld, V.G., *Hopf algebras and the quantum Yang-Baxter equation*, Soviet Math. Dokl. **32** (1985), 254-258.
- [FR] Frenkel, I.B., and Reshetikhin, N.Yu., *Quantum affine algebras and holonomic difference equations*, Comm. Math. Phys. **146** (1992), 1-60.
- [J1] Jimbo, M., *A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett.Math.Phys. **10** (1985), 63-69.
- [J2] Jimbo, M., *A q -analogue of $U(\mathfrak{gl}(N+1))$, Hecke algebra and the Yang-Baxter equation*, Lett. Math. Phys. **11** (1986), 247-252.
- [K] Karoubi, M., *K-Theory, An Introduction*, Springer-Verlag, New York, 1978.
- [Ka] Kashiwara, M., Comm. Math. Phys. (1990), 249-260.
- [KZ] Knizhnik, V.G., and Zamolodchikov, A.B., *Current algebra and Wess-Zumino model in two dimensions*, Nucl. Phys. **B247** (1984), 83-103.
- [M] Matsuo, A., *Jackson integrals of Jordan-Pochhammer type and quantum Knizhnik-Zamolodchikov equation*, Comm. Math. Phys. **to appear** (1992).
- [Mi] Milnor, J., and Stasheff, J., *Characteristic classes*, Princeton University Press, Princeton, New Jersey, U.S.A., 1974.
- [Mu] Mumford, D., *Projective invariants of projective structures and applications*, Proceedings of the International Congress of Mathematicians (1962), 526-530.
- [NS] Narasimhan, M.S., and Seshadri, C.S., *Stable and unitary vector bundles on a compact Riemann surface*, Ann.Math. **82** (1965), 540-567.
- [R] Reshetikhin, N.Yu., *Jackson-type integrals, Bethe vectors, and solutions to a difference analog of the Knizhnik-Zamolodchikov system*, preprint (1992).
- [SV] Schechtman, V.V., and Varchenko, A.N., *Arrangements of hyperplanes and Lie algebra homology*, Inv. Math. **106** (1991), 134-194.